

# SugarUp™

## 40% Glucose Gel

### Preservative-Free 40% Glucose Gel in Cups

#### SUGGESTED INSTRUCTIONS FOR USE

Single-patient use only.

Discard after opening.

Follow hospital protocol to determine dose and administration protocols.

Suggested protocol based on clinical literature:

1. Open container.
2. Draw up required dose of gel very slowly in an oral-only syringe.  
**Note: 3ml or 5ml syringes draw up gel more easily.**
3. Tap out any air bubbles.
4. Dry baby's mouth with gauze.
5. Dispense **one-half** dose from oral syringe onto gloved finger, in a stream - not a glob.
6. Massage into the buccal mucosa of one cheek.
7. Repeat with remaining half-dose inside the other cheek.
8. Large doses may be divided into four equal amounts and given alternating between cheeks.
9. Commence breast feeding, or administer expressed breast milk or formula by bottle.
10. Monitor infant according to hospital guideline.
11. **Note:** do not squirt gel directly into the baby's mouth. Always rub into the buccal mucosa where the blood flow insures absorption into the blood stream.

#### PRODUCT SPECIFICATIONS

Each cup contains 15ml of gel. Gel formulation: 40% dextrose (D-glucose), glycerine, USP purified water, maltodextrin, carboxymethyl cellulose, and citric acid (buffer). 12 month shelf life. Latex free, DEHP free, BPA free. Manufactured in the USA.

#### NEED A PROTOCOL FOR GLUCOSE?

Email us at [orders@sandboxmedical](mailto:orders@sandboxmedical) to request copies of protocols found in clinical literature or to share your protocol.

SugarUp™ is manufactured in the USA for SandBox Medical, LLC.



Cups may be recycled with other #5 plastics (polypropylene).

*SandBox Medical LLC*

750 Corporate Park Drive, Pembroke, MA 02359 USA  
Tel 877.396.1551 | 781.826.6905 Fax 781.826.8210  
[orders@sandboxmedical.com](mailto:orders@sandboxmedical.com) [sandboxmedical.com](http://sandboxmedical.com)

3-17021-003d

# SugarUp™

## 40% Glucose Gel

### Preservative-Free 40% Glucose Gel in Cups

#### SUGGESTED INSTRUCTIONS FOR USE

Single-patient use only.

Discard after opening.

Follow hospital protocol to determine dose and administration protocols.

Suggested protocol based on clinical literature:

1. Open container.
2. Draw up required dose of gel very slowly in an oral-only syringe.  
**Note: 3ml or 5ml syringes draw up gel more easily.**
3. Tap out any air bubbles.
4. Dry baby's mouth with gauze.
5. Dispense **one-half** dose from oral syringe onto gloved finger, in a stream - not a glob.
6. Massage into the buccal mucosa of one cheek.
7. Repeat with remaining half-dose inside the other cheek.
8. Large doses may be divided into four equal amounts and given alternating between cheeks.
9. Commence breast feeding, or administer expressed breast milk or formula by bottle.
10. Monitor infant according to hospital guideline.
11. **Note:** do not squirt gel directly into the baby's mouth. Always rub into the buccal mucosa where the blood flow insures absorption into the blood stream.

#### PRODUCT SPECIFICATIONS

Each cup contains 15ml of gel. Gel formulation: 40% dextrose (D-glucose), glycerine, USP purified water, maltodextrin, carboxymethyl cellulose, and citric acid (buffer). 12 month shelf life. Latex free, DEHP free, BPA free. Manufactured in the USA.

#### NEED A PROTOCOL FOR GLUCOSE?

Email us at [orders@sandboxmedical](mailto:orders@sandboxmedical) to request copies of protocols found in clinical literature or to share your protocol.

SugarUp™ is manufactured in the USA for SandBox Medical, LLC.



Cups may be recycled with other #5 plastics (polypropylene).

*SandBox Medical LLC*

750 Corporate Park Drive, Pembroke, MA 02359 USA  
Tel 877.396.1551 | 781.826.6905 Fax 781.826.8210  
[orders@sandboxmedical.com](mailto:orders@sandboxmedical.com) [sandboxmedical.com](http://sandboxmedical.com)

3-17021-003d