



40% Glucose Gel

Prefilled Oral-Only Syringes with Preservatives

SUGGESTED INSTRUCTIONS FOR USE



Follow your hospital protocol to determine weight specific dose and administration protocols.

Single-patient use only.
Discard after opening.

Suggested protocol based on clinical literature:

1. Open bag.
2. Tap out any air bubbles.
3. Dry baby's mouth with gauze.
4. Dispense **one-half** dose from oral syringe onto gloved finger, in a stream - not a glob. The gel is thick enough to stay on the finger.
5. Massage into the buccal mucosa of one cheek.
6. Administer remaining half-dose inside alternate cheek.
7. Large doses may be divided into four equal amounts and given alternating between cheeks.
8. Commence breast feeding, or administer expressed breast milk or formula by bottle.
9. Monitor infant according to hospital guideline.
10. **Note:** do not squirt gel directly into the baby's mouth. Always rub into the buccal mucosa where the blood flow insures absorption into the blood stream.

PRODUCT SPECIFICATIONS

Barcoded, individually bagged.
 Each 3ml oral-only syringe contains approximately 2.25ml of gel.
 Gel formulation: 40% dextrose (D-glucose), USP purified water, glycerine, maltodextrin, carboxymethyl cellulose, methylparaben & potassium sorbate (preservatives), citric acid & NaCl (sodium chloride) (buffers).
 24 month shelf life in unopened polybags. Latex free, DEHP free, BPA free.
 Gel made and packaged in the USA. Syringe made in China.
 Note: Preservatives are the same as used in SweetUms™ 24% Sucrose solution.

NEED A PROTOCOL FOR GLUCOSE?

Email us at orders@sandboxmedical to request copies of protocols found in clinical literature or to share your protocol.

SugarUp™ is manufactured in the USA for SandBox Medical, LLC.



Cups may be recycled with other #5 plastics (polypropylene).

SandBox Medical LLC

Tel 877.396.1551 | 781.826.6905 Fax 781.826.8210
orders@sandboxmedical.com sandboxmedical.com

3-17022-003b



40% Glucose Gel

Prefilled Oral-Only Syringes with Preservatives

SUGGESTED INSTRUCTIONS FOR USE



Follow your hospital protocol to determine weight specific dose and administration protocols.

Single-patient use only.
Discard after opening.

Suggested protocol based on clinical literature:

1. Open bag.
2. Tap out any air bubbles.
3. Dry baby's mouth with gauze.
4. Dispense **one-half** dose from oral syringe onto gloved finger, in a stream - not a glob. The gel is thick enough to stay on the finger.
5. Massage into the buccal mucosa of one cheek.
6. Administer remaining half-dose inside alternate cheek.
7. Large doses may be divided into four equal amounts and given alternating between cheeks.
8. Commence breast feeding, or administer expressed breast milk or formula by bottle.
9. Monitor infant according to hospital guideline.
10. **Note:** do not squirt gel directly into the baby's mouth. Always rub into the buccal mucosa where the blood flow insures absorption into the blood stream.

PRODUCT SPECIFICATIONS

Barcoded, individually bagged.
 Each 3ml oral-only syringe contains approximately 2.25ml of gel.
 Gel formulation: 40% dextrose (D-glucose), USP purified water, glycerine, maltodextrin, carboxymethyl cellulose, methylparaben & potassium sorbate (preservatives), citric acid & NaCl (sodium chloride) (buffers).
 24 month shelf life in unopened polybags. Latex free, DEHP free, BPA free.
 Gel made and packaged in the USA. Syringe made in China.
 Note: Preservatives are the same as used in SweetUms™ 24% Sucrose solution.

NEED A PROTOCOL FOR GLUCOSE?

Email us at orders@sandboxmedical to request copies of protocols found in clinical literature or to share your protocol.

SugarUp™ is manufactured in the USA for SandBox Medical, LLC.



Cups may be recycled with other #5 plastics (polypropylene).

SandBox Medical LLC

Tel 877.396.1551 | 781.826.6905 Fax 781.826.8210
orders@sandboxmedical.com sandboxmedical.com

3-17022-003b